

# High fructose consumption combined with low dietary magnesium intake may increase the incidence of the metabolic syndrome by inducing inflammation\*

Y. Rayssiguier<sup>1</sup>, E. Gueux<sup>1</sup>, W. Nowacki<sup>2</sup>, E. Rock<sup>1</sup>, A. Mazur<sup>1</sup>

<sup>1</sup> INRA, Unité de Nutrition Humaine, Clermont Ferrand/Theix, 63122 Saint-Genès-Champanelle, France; <sup>2</sup> Veterinary Faculty, Norwida Str. 31, 50375 Wrocław, Poland

**Correspondence:** Y. Rayssiguier, INRA, Unité de Nutrition Humaine, Clermont Ferrand – Theix, 63122 Saint-Genès-Champanelle, France <yrayssig@clermont.inra.fr>

**Abstract.** The metabolic syndrome is a cluster of common pathologies: abdominal obesity linked to an excess of visceral fat, insulin resistance, dyslipidemia and hypertension. This syndrome is occurring at epidemic rates, with dramatic consequences for human health worldwide, and appears to have emerged largely from changes in our diet and reduced physical activity. An important but not well-appreciated dietary change has been the substantial increase in fructose intake, which appears to be an important causative factor in the metabolic syndrome. There is also experimental and clinical evidence that the amount of magnesium in the western diet is insufficient to meet individual needs and that magnesium deficiency may contribute to insulin resistance. In recent years, several studies have been published that implicate subclinical chronic inflammation as an important pathogenic factor in the development of metabolic syndrome. Pro-inflammatory molecules produced by adipose tissue have been implicated in the development of insulin resistance. The present review will discuss experimental evidence showing that the metabolic syndrome, high fructose intake and low magnesium diet may all be linked to the inflammatory response. In many ways, fructose-fed rats display the changes observed in the metabolic syndrome and recent studies indicate that high-fructose feeding is associated with NADPH oxidase and renin-angiotensin activation. The production of reactive oxygen species results in the initiation and development of insulin resistance, hyperlipemia and high blood pressure in this model. In this rat model, a few days of experimental magnesium deficiency produces a clinical inflammatory syndrome characterized by leukocyte and macrophage activation, release of inflammatory cytokines, appearance of the acute phase proteins and excessive production of free radicals. Because magnesium acts as a natural calcium antagonist, the molecular basis for the inflammatory response is probably the result of a modulation of the intracellular calcium concentration. Potential mechanisms include the priming of phagocytic cells, the opening of calcium channels, activation of N-methyl-D-aspartate (NMDA) receptors, the activation of nuclear factor-kappaB (NFkB) and activation of the renin-angiotensin system. Since magnesium deficiency has a pro-inflammatory effect, the expected consequence would be an increased risk of developing insulin resistance when magnesium deficiency is combined with a high-fructose diet. Accordingly, magnesium deficiency combined with a high-fructose diet induces insulin resistance, hypertension, dyslipidemia, endothelial activation and prothrombic changes in combination with the upregulation of markers of inflammation and oxidative stress.

**Key words:** magnesium, fructose, metabolic syndrome, inflammation

The metabolic syndrome is a cluster of common pathologies, including abdominal obesity linked to an excess of visceral fat, insulin resistance, dyslipidemia and hypertension. This syndrome and type 2 diabetes are occurring at epidemic rates with frightful consequences for human health worldwide [1]. The increased prevalence of insulin resistance is linked to the western diet and reduced physical activity. In the past, diets high in saturated fats have been shown to induce insulin resistance and hyperlipidemia. However, recent studies suggest that a high intake of refined carbohydrates may also contribute to the risk of developing insulin resistance. In animal models, diets high in fructose have specifically been shown to contribute to a metabolic disturbance leading to insulin resistance. A significant change in the western diet is the substantial increase in dietary fructose consumption, which is due to a high intake of sucrose and high fructose corn syrup, a common sweetener used in the food industry [1]. Moreover, an increase in the consumption of empty calorie foods contributes to a decrease in the total mineral and vitamin density per unit of energy consumed and consequently, the marginal magnesium intake induces a high prevalence of marginal magnesium deficiency. A healthy magnesium intake supports efficient insulin function, whereas magnesium deficiency contributes to insulin resistance [2-6]. In industrialized countries, magnesium intake has been reduced, while fructose consumption has been rapidly increasing, and the aim of this review is to emphasize the consequences of this eating pattern, particularly in the development of the metabolic syndrome and to discuss the pathogenic role of inflammation.

### **Role of inflammation in the metabolic syndrome**

In recent years, several studies have demonstrated that subclinical chronic inflammation is an important pathogenic factor in the development of insulin resistance and cardiovascular diseases [7, 8]. Markers for this inflammatory response include acute phase proteins, cytokines and mediators associated with endothelial activation. In obese individuals, the white adipose tissue contains an increased number of macrophages, when compared with lean individuals, and the macrophages appear to be activated. The adipocytes and macrophages produce leptin and other factors, which upregulate the number of adhesion molecules on endothelial cells. This leads to transmigration of monocytes, an increase in white

adipose tissue-resident macrophages and consequently, cytokines are released in large amounts from the adipose tissue [9]. According to this hypothesis, an excess energy intake leads to obesity and hyperglycemia, which can cause oxidative stress and inflammatory changes (nuclear factor-kappaB [NFkB] activation, increased levels of tumour necrosis factor [TNF] alpha and interleukin [IL] [6]). These inflammatory changes inhibit insulin signalling and can lead to insulin resistance. Moreover, the inflammatory state induces beta cells dysfunction, which in combination with insulin resistance leads to type 2 diabetes [10, 11].

### **High-fructose diet and metabolic syndrome**

The fructose fed rat represents a model for the metabolic syndrome, including insulin resistance, hypertension and dyslipidemia, and growing evidence suggests a role for inflammation and oxidative stress in this model [12]. High fructose feeding of rats is associated with a rapid increase in reactive oxygen species (ROS) production by polymorphonuclear leukocytes (PMN) and tissues. This overproduction of ROS is dependent on NADPH oxidase activation and leads to an oxidative stress response as evidenced by elevated concentrations of plasma and urinary thiobarbituric acid reactive substances (TBARS). In rats, the fructose diet, when compared with a starch diet, lowers the plasma vitamin E/triglycerides (TG) ratio, which results in an increase in TG-rich lipoprotein susceptibility to lipid peroxidation, and hearts are less protected against *in vitro* peroxidation [13, 14]. Recent findings suggest that the pro-oxidant effect induced by a high fructose diet can be decreased by consumption of fermentable carbohydrates such as oligofructose [15]. The link between inflammation, oxidative stress and metabolic syndrome is supported by the fact that females, which are protected against the pro-oxidant effects of a high-fructose diet when compared with males, do not develop insulin resistance [16]. Diets rich in fructose can alter cellular metabolism via several pathways, thereby accelerating oxidative stress. Fructose feeding results in the activation of the renin-angiotensin system and it is well-established that angiotensin 2 production is associated with oxidative stress. This oxidative stress is characterized by overproduction of ROS and is dependent on the activation of NADPH oxidase [17]. Moreover, increased IL1 and IL6 plasma levels have been documented in rats fed a high-fructose diet [18]. Finally, in addition to the possible roles of uric acid and kidney damage

[19], the inflammatory response and oxidative stress may be key events in the development of insulin resistance in rats fed a high fructose diet. These events are likely to contribute to other aspects of the metabolic syndrome. The hypertriglyceridemic effect of fructose can be reduced by lipooxygenase inhibitors, which decrease the inflammatory response [20], and inflammation and oxidative stress are important factors in cardiovascular effects.

#### **Inflammatory effects of magnesium deficiency**

A characteristic allergy-like crisis occurs spontaneously in magnesium deficient rats. They present hyperemia and oedema of the ears and legs, and a greater spleen size due to phagocytic cell infiltration [5, 6, 21]. An inflammatory response has also been observed in other magnesium-deficient rodents. However, it was shown that in mice the response depends on the experimental conditions [22-24]. The inflammatory syndrome is accompanied by hyperalgesia, which can be prevented by an N-methyl-D-aspartate (NMDA) receptor antagonist [25]. In blood, the most prominent change is leukocytosis, which results from the increased number of PMN leukocytes, mainly neutrophils and eosinophils. An increase in the number of eosinophils is considered a typical allergic response. An important phenomenon during inflammation is the production of cytokines. Early studies reported that, in rats, experimental magnesium deficiency led to increased plasma levels of orosomucoid [26]. In the same experimental model, several positive acute phase protein plasma concentrations, and/or their liver mRNA levels, have been shown to be induced, including alpha2-macroglobulin, alpha1-acid glycoprotein, complement, fetoprotein, haptoglobin and fibrinogen. These changes are related to increased IL6 concentrations, which stimulate the synthesis of many acute phase proteins by the liver. The decrease of several negative acute phase proteins, such as albumin, apolipoprotein (apo) E, retinol-binding protein (RBP), is a classical finding of an acute phase response. This inflammatory response and its consequences are only observed in male rats. Female animals are partially protected, suggesting that estrogens may be protective against the proinflammatory effects of magnesium deficiency [27]. It is of particular interest that females, when compared with males, do not develop the metabolic syndrome on the fructose diet. Several studies have been performed to assess the activation of inflammatory cells in magnesium deficiency [28-30]. Free radical generation from PMN

was measured *in vitro* using chemiluminescence, and via NADPH oxidase activation, both macrophages and neutrophils generated superoxide anions in response to various stimuli. There was a low basal neutrophil activity in control rats, but the basal neutrophil activity of magnesium deficient rats was significantly higher. Neutrophils from control and magnesium-deficient rats were responsive to activation by phorbol 12-myristate 13-acetate (PMA) and zymosan, and the response was higher for PMN magnesium-deficient rats. Differential gene expression analysis of stress proteins has confirmed the neutrophil activation [30] and in fact, the majority of stress proteins were upregulated in neutrophils from magnesium-deficient animals. Resident macrophages of magnesium-deficient rats present the morphological aspects of an activated cell and their chemiluminescence activity is elevated. *In vitro*, these cells are more sensitive to PMA stimulation. Magnesium-deficient rats are more sensitive to immune stress as measured by TNF response following an endotoxin challenge [31]. The specific mechanisms of the inflammatory response in magnesium deficiency have not been elucidated. However, magnesium acts as a natural calcium antagonist and the molecular basis for the inflammatory response is probably linked to the modulation of the intracellular calcium concentration [32]. Potential mechanisms include the priming of phagocytic cells, the opening calcium channels and activation of NMDA receptors [33], the release of neurotransmitters such as substance P [34, 35], the activation of NFkB [36] and the activation of the renin angiotensin systems [37].

#### **High-fructose diet combined with magnesium deficiency**

Inflammation is a key event in the initiation and development of metabolic syndrome. A high-fructose diet induces both inflammation and the metabolic syndrome in rats. If magnesium deficiency has a proinflammatory effect, one likely consequence of combining magnesium deficiency with a high-fructose diet is an amplified effect, which has been shown by recent data from rats. The fructose diet, as compared to a starch diet, induces hyperglycemia and hyperinsulinemia. Magnesium deficiency induces hyperglycemia and hyperinsulinemia when compared to a control diet. The combined effects are higher, using the high fructose low magnesium diet, and the binding of insulin to the red blood cell (RBC) insulin receptors is reduced [38]. Thus, this study clearly documents the effects of magnesium defi-

ciency in the development of insulin resistance in the rat model. We have shown that fructose feeding in magnesium deficient rats is associated with an increase in oxidative stress when compared with magnesium deficient rats fed a starch diet [39]. This observation is consistent with the hypothesis that inflammation and increased oxidative stress, induced by magnesium deficiency, contribute to the development of insulin resistance. In the same model, the proinflammatory effect of magnesium deficiency contributes to other aspects of the metabolic syndrome: hyperlipemia, elevated blood pressure, endothelial dysfunction and increased thrombosis tendency [5, 6]. Fructose has been shown to increase the hyperlipemic effect of magnesium deficiency, when compared to starch [40]. As previously discussed, magnesium deficiency is accompanied by alterations in lipid metabolism including accumulation of TG-rich lipoproteins (TGRLP), a decrease in high density lipoprotein (HDL) levels, an increase in apoB, a decrease in apoA1 and apoE, modifications in the composition of lipoproteins, and a defect in the clearance of TGRLP [41, 42]. Several factors contribute to these alterations including decreases in the activities of lipoprotein lipase, hepatic lipase [43, 44] and lecithin-cholesterol acyl transferase (LCAT) [45]. Together, these changes could contribute to decreased reverse cholesterol transport. TGRLP and tissues from magnesium-deficient rats are more susceptible to *ex vivo* oxidation than those from controls [46-48]. Thus, oxidative modification of lipoproteins could play a significant role in the pathogenesis of vascular lesions following magnesium deficiency and furthermore, magnesium affects the inflammatory-dependent events leading to atherosclerosis. The lipid metabolism changes observed in experimental magnesium deficiency have been observed in other models in inflammation [49]. On the basis of these studies, it is possible to conclude that the proatherogenic lipoprotein changes in magnesium deficient rats are the consequence of the inflammatory response. Epidemiological and experimental studies have demonstrated an inverse association between magnesium status and blood pressure [50, 51]. Chronic dietary magnesium deficiency causes elevated blood pressure; initially, a hypotension phase is observed, which is due to the release of inflammatory agents, the subsequent hypertension is a result of oxidative stress and structural modifications in the vascular system [52, 53]. In fact, free radicals may partly inactivate NO and an increased degradation of NO by superoxide anions could contribute to hypertension during chronic magnesium deficiency. *In vitro* studies have shown that low

magnesium levels results in endothelial dysfunction [54, 55]. Serum from magnesium deficient rats stimulates the proliferation of endothelial cells, increases the adhesion of monocytes to these cells, and the endothelial cells upregulate the plasminogen activator inhibitor (PAI) factor. The presence of endothelial dysfunction and dyslipidemia triggers platelet aggregability [56]. Platelet aggregation is increased in magnesium deficiency, while magnesium supplementation inhibits platelet dependent thrombosis [44, 57]. We have investigated the thrombotic tendency in rats fed a magnesium-deficient, high fructose diet. Following epinephrine, the magnesium-deficient rats exhibited lesions in left atrium which was dilated and haemorrhagic [44]. Thus, magnesium deficiency induces predisposition to epinephrine-initiated thrombosis and inflammatory pathway promotes thrombosis which is responsible for myocardial infarction and stroke [28, 44].

## Conclusion

Magnesium deficiency combined with a high-fructose diet induces insulin resistance, hypertension, dyslipidemia, endothelial activation and prothrombotic changes. In addition to well-known factors such as overnutrition and physical inactivity, a diet rich in fructose and deficient in magnesium results in an inflammatory response, which may represent a triggering factor in the development of the metabolic syndrome. Our data are in good agreement with clinical data [58-66] showing that low serum and dietary magnesium levels are strongly correlated with low grade systemic inflammation and metabolic syndrome [58-66]. Large epidemiological studies indicate that lower serum magnesium levels are associated with insulin resistance. Magnesium intake and systemic inflammation and the prevalence of metabolic syndrome were inversely correlated in subjects participating in the Women Health Study [62]. Furthermore, low magnesium serum levels are associated with elevated serum concentrations of both TNF-alpha [64] and C-reactive protein (CRP) [62, 63], suggesting that magnesium deficiency may also be involved in the development of the low-grade chronic inflammatory syndrome which can induce metabolic disorders. Other studies have shown the correlation between low-serum magnesium levels and TNF-alpha in obese and non-alcoholic steatohepatitis (NASH) subjects [65]. Moreover, a longitudinal association of magnesium intake with the incidence of metabolic syndrome has recently been examined and the data suggests that

young adults with higher magnesium intake have a lower risk of developing the metabolic syndrome [66]. A typical western diet is high in fructose and often magnesium deficient. Considering the well-documented detrimental effect of this eating pattern, there is urgent need to explore the details of the link between diet and the inflammatory signalling pathways.

## References

1. Basciano H, Federico L, Adeli K. Fructose, insulin resistance, and metabolic dyslipidemia. *Nutr Metab* 2005; 2: 5.
2. Lopez-Ridaura R, Willett WC, Rimm EB, Liu S, Stampfer MJ, Manson JE, Hu FB. Magnesium intake and risk of type 2 diabetes in men and women. *Diabetes Care* 2004; 27: 134-40.
3. Tayaka J, Higashino H, Kobayashi Y. Intracellular magnesium and insulin resistance. *Magnes Res* 2004; 17: 126-36.
4. Ueshima K. Magnesium and ischemic heart disease: a review of epidemiological, experimental, and clinical evidences. *Magnes Res* 2005; 18: 275-84.
5. Rayssiguier Y, Mazur A. Magnesium and inflammation: lessons from animal models. *Clin Calcium* 2005; 15: 101-4.
6. Mazur A, Maier JAM, Rock E, Gueux E, Nowacki W, Rayssiguier Y. Magnesium and the inflammatory response: potential pathophysiological implications. *Arch Biochem Biophys* 2006; (in press).
7. Dandona P, Aljada A, Bandyopadhyay A. Inflammation: the link between insulin resistance, obesity and diabetes. *Trends Immunol* 2004; 25: 4-7.
8. Fantuzzi G. Adipose tissue, adipokines, and inflammation. *J Allergy Clin Imm* 2005; 115: 911-9.
9. Rudin E, Barzilai N. Inflammatory peptides derived from adipose tissue. *Immun Ageing* 2005; 2: 1-3.
10. Sjöholm A, Nyström T. Endothelial inflammation in insulin resistance. *Lancet* 2005; 365: 610-2.
11. Sjöholm A, Nyström T. Inflammation and the etiology of type 2 diabetes. *Diabetes Metab Res Rev* 2006; 22: 4-10.
12. Delbosc S, Paizanis E, Magous R, Araiz C, Dimo T, Cristo JP, Cros G, Azay Y. Involvement of oxidative stress and NADPH oxidase activation in the development of cardiovascular complications in a model of insulin resistance, the fructose-fed rat. *Atherosclerosis* 2005; 179: 43-9.
13. Busserolles J, Rock E, Gueux E, Mazur A, Grolier P, Rayssiguier Y. Short-term consumption of a high-sucrose diet has a pro-oxidant effect in rats. *Br J Nutr* 2002; 87: 337-42.
14. Busserolles J, Zimowska W, Rock E, Rayssiguier Y, Mazur A. Rats fed a high sucrose diet have altered heart antioxidant enzyme activity and gene express. *Life Sci* 2002; 71: 1303-12.
15. Busserolles J, Gueux E, Rock E, Demigne C, Mazur A, Rayssiguier Y. Oligofructose protects against the hypertriglyceridemic and pro-oxidative effects of a high fructose diet in rats. *J Nutr* 2003; 133: 1903-8.
16. Busserolles J, Mazur A, Gueux E, Rock E, Rayssiguier Y. Metabolic syndrome in the rats: females are protected against the pro-oxidant effect of a high sucrose diet. *Exp Biol Med* 2002; 227: 837-42.
17. Shinozaki K, Ayajiki K, Nishio Y, Sugaya T, Kashiwagi A, Okamura T. Evidence for a causal role of the renin-angiotensin system in vascular dysfunction associated with insulin resistance. *Hypertension* 2004; 43: 255-62.
18. Armutcu F, Coskun O, Gurel A, Kanter M, Can M, Ucar F, Unalacak M. Thymosin alpha 1 attenuates lipid peroxidation and improves fructose-induced steatohepatitis in rats. *Clin Biochem* 2005; 38: 540-7.
19. Nakagawa T, Hu H, Zharikov S, Tuttle KR, Short RA, Glushakova O, Ouyang X, Feig DI, Block ER, Herrera-Acosta J, Patel JM, Johnson RJ. A causal role for uric acid in fructose-induced metabolic syndrome. *Am J Physiol Renal Physiol* 2006; 290: 625-31.
20. Kelley GL, Azhar S. Reversal of high dietary fructose-induced PPARalpha suppression by oral administration of lipoxygenase/cyclooxygenase inhibitors. *Nutr Metab (Lond)* 2005; 2: 18.
21. Malpuech-Brugère C, Nowacki W, Daveau M, Gueux E, Linard C, Rock E, Lebreton J, Mazur A, Rayssiguier Y. Inflammatory response following acute magnesium deficiency in the rat. *Biochim Biophys Acta* 2000; 1501: 91-8.
22. Nasulewicz A, Zimowska W, Bayle D, Dzimira S, Madej J, Rayssiguier Y, Opolski A, Mazur A. Changes in gene expression in the lungs of Mg-deficient mice are related to an inflammatory process. *Magnes Res* 2004; 17: 259-63.
23. Sabbagh F, Lecerf F, Maurois P, Bac P, German-Fattal M. Severe Mg-deficiency is not associated with endothelial cell activation in mouse lung. *Magnes Res* 2005; 18: 225-34.
24. Sabbagh F, Lecerf F, Maurois P, Bac P, German-Fattal M. Effect of Mg-deficiency on endothelial cell allogenic activation in a model of isolated perfused mouse lung. *Magnes Res* 2005; 18: 235-40.
25. Dubray C, Alloui A, Bardin L, Rock E, Mazur A, Rayssiguier Y, Eschalier A, Lavarenne J. 1997; Magnesium deficiency induces an hyperalgesia reversed by the NMDA receptor antagonist MK801. *Neuroreport* 1997; 8: 1383-6.
26. Augusti Y, Rayssiguier Y, Larvor P, Durlach J. Magnesium deficiency in rats and variation of certain serum protein parameters. *C R Seances Soc Biol Fil* 1973; 167: 1804-8.
27. Bussièrè F, Gueux E, Rock E, Mazur A, Rayssiguier Y. Female rats are less susceptible to inflammation induced by magnesium deficiency than males: the influence of estrogen. In: Rayssiguier Y, Mazur A, Durlach J, eds. *Advances in Magnesium Research: Nutrition and Health*. John Libbey & Company Ltd, 2001: 313-4.

28. Rayssiguier Y, Bussi re F, Gueux E, Rock E, Mazur A. Acute phase response in magnesium deficiency: possible relevance to atherosclerosis. In: Rayssiguier Y, Mazur A, Durlach J, eds. *Advances in Magnesium Research: Nutrition and Health*. John Libbey & Company Ltd, 2001: 277-83.
29. Bussi re F, Gueux E, Rock E, Girardeau JP, Tridon A, Mazur A, Rayssiguier Y. Increased phagocytosis and production of reactive oxygen species by neutrophils during magnesium deficiency in rats and inhibition by high magnesium concentration. *Br J Nutr* 2002; 87: 107-13.
30. Bussi re FI, Zimowska W, Gueux E, Rayssiguier Y, Mazur A. Stress protein expression cDNA array study supports activation of neutrophils during acute magnesium deficiency in rats. *Magnes Res* 2002; 15: 37-42.
31. Malpuech-Brug re C, Rock E, Astier C, Nowacki W, Mazur A, Rayssiguier Y. Exacerbated immun stress response during experimental magnesium deficiency results from abnormal cell calcium homeostasis. *Life Sci* 1998; 63: 1815-22.
32. Bussi re FI, Gueux E, Rock E, Mazur A, Rayssiguier Y. Protective effect of calcium deficiency on the inflammatory response in magnesium-deficient rats. *Eur J Nutr* 2002; 41: 197-202.
33. Tejero-Taldo I, Chmielinska JJ, Gonzalez G, Tong Mak I, Weglicki WB. N-Methyl-D-aspartate receptor blockade inhibits cardiac inflammation in the Mg<sup>2+</sup>-deficient rat. *JPET* 2004; 311: 1-8.
34. Weglicki WB, Mak IT, Kramer JH, Dickens BF, Cassidy MM, Stafford RE, Phillips TM. Role of free radicals and substance P in magnesium deficiency. *Cardiovasc Res* 1996; 31: 677-82.
35. O'Connor TM, O'Connell J, O'Brien DI, Goode T, Bredin CP, Shanahan F. The role of substance P in inflammatory disease. *J Cell Physiol* 2004; 201: 167-80.
36. Altura BM, Gebrewold A, Zhang A, Altura BT. Low extracellular magnesium ions induce lipid peroxidation and activation of nuclear factor-kappa B in canine cerebral vascular smooth muscle: possible relation to traumatic brain injury and strokes. *Neurosci Lett* 2003; 341: 189-92.
37. Sapna S, Ranjith SK, Shivakumar K. Cardiac fibrogenesis in magnesium deficiency: a role for circulating angiotensin II and aldosterone. *Am J Physiol Heart Circ Physiol* 2006; 291: H436-H440.
38. Chaudhary DP, Boparai RK, Sharma R, Bansal DD. Studies on the development of an insulin resistant rat model by chronic feeding of low magnesium high sucrose diet. *Magnes Res* 2004; 17: 293-300.
39. Busserolles J, Gueux E, Rock E, Mazur A, Rayssiguier Y. High fructose feeding of magnesium deficient rats is associated with increased plasma triglyceride concentration and increased oxidative stress. *Magnes Res* 2003; 16: 7-12.
40. Rayssiguier Y, Gueux E, Weiser D. Effect of magnesium deficiency on lipid metabolism in rats fed a high carbohydrate diet. *J Nutr* 1981; 111: 1876-83.
41. Nassir F, Mazur A, Giannoni F, Gueux E, Davidson NO, Rayssiguier Y. Magnesium deficiency modulates hepatic lipogenesis and apolipoprotein gene expression in the rat. *Biochim Biophys Acta* 1995; 1257: 125-32.
42. Gueux E, Mazur A, Cardot P, Rayssiguier Y. Magnesium deficiency affects plasma lipoprotein composition in rats. *J Nutr* 1991; 121: 1222-7.
43. Rayssiguier Y, Noe L, Etienne J, Gueux E, Cardot P, Mazur A. Effect of magnesium deficiency on post-heparin lipase activity and tissue lipoprotein lipase in the rat. *Lipids* 1991; 16: 182-6.
44. Rayssiguier Y, Gueux E. Magnesium and lipids in cardiovascular disease. *J Am Coll Nutr* 1986; 5: 507-19.
45. Gueux E, Rayssiguier Y, Piot MC, Alcindor L. Reduction of plasma lecithin-cholesterol acyltransferase activity by acute magnesium deficiency in the rat. *J Nutr* 1984; 114: 1479-83.
46. Rayssiguier Y, Gueux E, Bussi re L, Durlach J, Mazur A. Dietary magnesium affects susceptibility of lipoproteins and tissues to peroxidation in rats. *J Am Coll Nutr* 1993; 12: 133-7.
47. Bussi re L, Mazur A, Gueux E, Nowacki W, Rayssiguier Y. Triglyceride-rich lipoproteins from magnesium-deficient rats are more susceptible to oxidation by cells and promote proliferation of cultured vascular smooth muscle cells. *Magnes Res* 1995; 8: 151-7.
48. Gueux E, Azais-Braesco V, Bussi re L, Grolier P, Mazur A, Rayssiguier Y. Effect of magnesium deficiency on triacylglycerol-rich lipoprotein and tissue susceptibility to peroxidation in relation to vitamin E content. *Br J Nutr* 1995; 74: 849-56.
49. Khovidhunkit W, Memon RA, Feingold KR, Grunfeld C. Infection and inflammation-induced proatherogenic changes of lipoproteins. *J Infect Dis* 2000; 181: S462-S472.
50. Maier JAM. Low magnesium and atherosclerosis: an evidence-based link. *Mol Aspects Med* 2003; 24: 137-46.
51. Touyz RM. Role of magnesium in the pathogenesis of hypertension. *Mol Aspects Med* 2003; 24: 107-36.
52. Rayssiguier Y, Mbega JD, Durlach V, Gueux E, Durlach J, Giry J, Dalle M, Mazur A, Laurant P, Berthelot A. Magnesium and blood pressure. I. Animal studies. *Magnes Res* 1992; 5: 139-46.
53. Bukoski RD. Reactive oxygen species: the missing link between magnesium deficiency and hypertension? *J Hypertens* 2002; 20: 2141-3.
54. Maier JAM, Malpuech-Brugere C, Zimowska W, Rayssiguier Y, Mazur A. Low magnesium promotes endothelial cell dysfunction: implications for atherosclerosis, inflammation and thrombosis. *Biochim Biophys Acta* 2004; 1689: 13-21.
55. Bernardini D, Nasulewicz A, Mazur A, Maier JAM. Magnesium and microvascular endothelial cells: a role in inflammation and angiogenesis. *Front Biosc* 2005; 10.
56. Nieuwdorp M, Stroes ESG, Meijers JCM, B ller H. Hypercoagulability in the metabolic syndrome. *Curr Opin Pharmacol* 2005; 5: 155-9.

57. Shechter M, Merz CN, Paul-Labrador M, Meisel SR, Rude RK, Molloy MD, Dwyer JH, Shah PK, Kaul S. Oral magnesium supplementation inhibits platelet-dependent thrombosis in patients with coronary artery disease. *Am J Cardiol* 1999; 84: 152-6.
58. Guerrero-Romero F, Rodriguez-Moran M. Low serum magnesium levels and metabolic syndrome. *Acta Diabetol* 2002; 39: 209-13.
59. Barbagallo M, Dominguez LJ, Galioto A, Ferlisi A, Cani C, Malfa L, Pineo A, Busardo A, Paolisso G. Role of magnesium in insulin action, diabetes and cardio-metabolic syndrome X. *Mol Aspects Med* 2003; 24: 39-52.
60. Huerta MG, Roemmich JN, Kington ML, Bovbjerg VE, Weltman AL, Holmes VF, Patrie JT, Rogol AD, Nadler JL. Magnesium deficiency is associated with insulin resistance in obese children. *Diabetes Care* 2005; 28: 1175-81.
61. Guerrero-Romero F, Rodriguez-Moran M. Hypomagnesemia, oxidative stress, inflammation, and metabolic syndrome. *Diabetes Metab Res Rev* 2006; (in press).
62. Song Y, Ridker PM, Manson JE, Cook NR, Buring JE, Liu S. Magnesium intake, C-reactive protein, and the prevalence of metabolic syndrome in middle-aged and older U.S. women. *Diabetes Care* 2005; 28: 1438-44.
63. King DE, Mainous 3rd AG, Geesey ME, Woolson RF. Dietary magnesium and C-reactive protein levels. *J Am Coll Nutr* 2005; 24: 166-71.
64. Rodriguez-Morand M, Guerrero-Romero F. Elevated concentrations of TNF-alpha are related to low serum magnesium levels in obese subjects. *Magnes Res* 2004; 17: 189-96.
65. Rodriguez-Hernandez H, Gonzalez JL, Rodriguez-Moran M, Guerrero-Romero F. Hypomagnesemia, insulin resistance, and non-alcoholic steatohepatitis in obese subjects. *Arch Med Res* 2005; 36: 362-6.
66. He K, Liu K, Daviglus ML, Morris SJ, Loria CM, Van Horn L, Jacobs Jr. DR, Savage PJ. Magnesium intake and incidence of metabolic syndrome among young adults. *Circulation* 2006; 113: 1675-82.